2018 | Volume Volume - 3 - Issue Issue - 1

In this issue

Research Article

Open Access Research Article PTZAID:IJRRO-3-124

Cardiac Computed Tomography Coronary Angiography Post Non-diagnostic or **Equivocal Myocardial Perfusion SPECT**

Published On: May 31, 2017 | Pages: 015 - 020

Author(s): Abdullah Al-harbi, Mervat M Aboulkheir and Ahmed Fathala*

Background: Myocardial perfusion single- photon emission computed tomography (MP-SPECT) plays a key role in the management of patients with known or suspected coronary artery disease (CAD). However, Infrequently, MP-SPECT is non-diagnostic or equivocal. ...

Abstract View Full Article View DOI: 10.17352/ijrro.000024

Case Report

Open Access Case Report PTZAID:IJRRO-3-125

Radiation Recall Masquerading as an Infectious Process

Published On: June 29, 2017 | Pages: 021 - 024

Author(s): Santhosshi Narayanan, Bilal Mujtaba*, Bilal Mujtaba, Mohamed Elshikh, Mohamed Elshikh and Gauri R Varadhachary Cellulitis and other infections are common in cancer patients, especially patients who are immunosuppressed following chemotherapy. When patients do not respond to antimicrobial therapy as expected, it is imperative for clinicians to consider other treatment-related conditions in the differential diagnosis. ...

Abstract View Full Article View DOI: 10.17352/ijrro.000025

Open Access Case Report PTZAID:IJRRO-3-123

Giant Fibrolipoma Mimicking Subcutaneous Liposarcoma in Shoulder

Published On: February 01, 2017 | Pages: 013 - 014

Author(s): Özkan Özen, Abdullah Çelik, Alptekin Tosun*, Kürad Aytekin and Cem Zeki Esenyel

Lipomas are the most common lesions in subcutaneous tissues. Fibrolipomas are a rare subtype of lipomas. A patient with the mass growth on the left shoulder within 1 year applied to hospital and a large sized fi xed mass was detected on the localization. ...

Abstract View Full Article View DOI: 10.17352/ijrro.000023

Open Access Case Report PTZAID:IJRRO-3-122

Acoustic Radiation Force Impulse Imaging in the Differential Diagnosis of an **Ectopic Intrathyroidal Thymus**

Published On: January 27, 2017 | Pages: 010 - 012

Author(s): Dilek Salam*, Meltem Ceyhan Bilgici, Cengiz Kara, Muzaffer Elmali, Asli Tanrvermi Sayit

Ectopic intrathyroidal thymus is caused by a hesitate in the migration of thymus during embryogenesis and is a rare cause of a nodular intrathyroidal lesion. Nodular thyroid lesions raise the suspicion of malignancy in the pediatric age group. ...

Abstract View Full Article View DOI: 10.17352/iirro.000022

Open Access Case Report PTZAID:IJRRO-3-121

Contribution of Tc-99m RBC SPECT/ CT in Intrahepatic and Peritoneal Splenosis Mimicking Malignancy

Published On: January 25, 2017 | Pages: 007 - 009

Author(s): Kevser Oksuzoglu*, Rabia Ergelen, Tunc Ones, Sabahat Inanir and Tanju Yusuf Erdil

Splenosis is the autoimplantation of ectopic spleen tissue resulting heterotopic auto-transplantation and implantation of splenic tissue after splenic ruptur caused by trauma or splenectomy. ...

Abstract View Full Article View DOI: 10.17352/ijrro.000021

Open Access Case Report PTZAID:IJRRO-3-120

Combination of Thrombolysis and Glycoprotein Ilb/Illa Inhibition in Chronic **Peripheral Thrombosis: A Case Report**

Published On: January 24, 2017 | Pages: 004 - 006

Author(s): Mubin I Syed*, Ryan Gallagher, Zeshan Hussain, Azim Shaikh, Philip J Cain, Kamal Morar, Matthew Sebastian, Robert Tyrrell, Allen Guehl and Ronald Erdelyi

Currently, many interventionalists employ either tissue plasminogen activator (alteplase, Genentech, San Francisco, CA), or recombinant plasminogen activator (reteplase, Chiesi USA, Inc., Cary, NC), for peripheral thrombolysis. ...

Abstract View Full Article View DOI: 10.17352/ijrro.000020

Open Access Case Report PTZAID:IJRRO-3-119

Adenoid Cystic Carcinoma of the Maxilla, CT Follow-Up: A Case Report

Published On: January 07, 2017 | Pages: 001 - 003

Author(s): Bruno Vieira Caputo*, Gilberto Araujo Noro Filho, Claudio Costa, Artur Aburad de Carvalhosa and Elcio Magdalena Giovani

Background: Intraoral adenoid cystic carcinoma is an uncommon pathology with a variable clinical course, it develops as a slow growing swelling characterized by wide local infiltration. Most patients are in their fourth and sixth decades of life, and females are slightly more affected than males. ...

Abstract View Full Article View DOI: 10.17352/ijrro.000019